

ITP 93-9
May 1993

FORM CONNECTIONS

Ingemar Bengtsson

Abstract

Riemannian geometry in four dimensions naturally leads to an $SL(3)$ connection that annihilates a basis for self-dual two-forms. Einstein's equations may be written in terms of an $SO(3)$ connection, with $SO(3)$ chosen as an appropriate subgroup of $SL(3)$. We show how a set of "neighbours" of Einstein's equations arises because the subgroup may be chosen in different ways. An explicit example of a non-Einstein metric obtained in this way is given. Some remarks on three dimensional space-times are made.

Institute of Theoretical Physics
S-41296 Göteborg
Sweden

As is by now well known, it is possible to formulate Einstein's equations using a configuration space that is spanned by an $\text{SO}(3)$ connection [1] [2]. In a related development, 'tHooft [3] found an action for the same equations which employs an $\text{SL}(3)$ connection as the basic variable. The purpose of this letter is to use a part of the latter formalism to illuminate some aspects of the former, in particular the surprising possibility [4] to construct diffeomorphism invariant theories which do not lead to Einstein geometries, even though they have the same number of degrees of freedom as Einstein's theory. Please note that many of the ideas which we will use have been around for a long time [5].

Although we will later make some comments on three-dimensional spaces, most of our results are peculiar to four dimensions, and they all hinge on the fact that the six-dimensional space $\mathbf{W}$ of two-forms on a four-dimensional vector space may be split into two orthogonal subspaces, with a scalar product given by

$$(\Sigma_1, \Sigma_2) \equiv \frac{1}{2} \epsilon^{\alpha\beta\gamma\delta} \Sigma_{\alpha\beta 1} \Sigma_{\gamma\delta 2}. \quad (1)$$

Moreover it turns out that there is a one-to-one correspondence between all the ways in which $\mathbf{W}$ may be split into two orthogonal subspaces $\mathbf{W}^+$ and $\mathbf{W}^-$ and the space of conformal structures on the original vector space. Assuming that the triplet of two-forms $\Sigma_{\alpha\beta i}$ forms a basis for $\mathbf{W}^+$, it may be shown that these basis vectors are by construction self-dual with respect to the metric

$$g_{\alpha\beta} \equiv \frac{8}{3} \sigma \epsilon^{ijk} \Sigma_{\alpha\gamma i} \tilde{\Sigma}_j^{\gamma\delta} \Sigma_{\delta\beta k} \quad (2)$$

where the conformal factor σ is so far arbitrary, the volume element becomes

$$\sqrt{-g} = \frac{i}{2} \sigma^2 \det m, \quad (3)$$

and we use the notation

$$\tilde{\Sigma}^{\alpha\beta} \equiv \frac{1}{2} \epsilon^{\alpha\beta\gamma\delta} \Sigma_{\gamma\delta} \quad (4)$$

$$m_{ij} \equiv \epsilon^{\alpha\beta\gamma\delta} \Sigma_{\alpha\beta i} \Sigma_{\gamma\delta j}. \quad (5)$$

The proof of this statement [6] is an elegant application of Clifford algebra ideas, and starts from the elementary observation that

$$\Sigma_{\alpha\gamma i}\tilde{\Sigma}_j^{\gamma\beta} + \Sigma_{\alpha\gamma j}\tilde{\Sigma}_i^{\gamma\beta} = -1/4m_{ij}\delta_\alpha^\beta. \quad (6)$$

(In eq. (6) only, the index i is not restricted to run from one to three. This should cause no confusion.)

Note that we have chosen conventions appropriate to the case of Lorentzian signatures, although it should be said that the conditions that guarantee that we obtain a real metric with a particular signature is so far fully understood (in the context of the field equations that we will present) only in the Euclidean and neutral cases.

With these facts, and the assumption that the Σ 's are non-degenerate in the sense that they may serve as a basis for the self-dual subspace, a pair of very useful formulas follows:

$$\Sigma_\alpha{}^\gamma{}_i \Sigma_\gamma{}^\beta{}_j = \frac{i}{8\sqrt{-g}} m_{ij} \delta_\alpha^\beta - \frac{1}{2} \sigma^{-1} \epsilon_{ijk} m^{km} \Sigma_\alpha{}^\beta{}_m \quad (7)$$

$$\Sigma_{\alpha\beta i} m^{ij} \Sigma_{\gamma\delta j} = \frac{1}{8} \epsilon_{\alpha\beta\gamma\delta} - \frac{i}{8\sqrt{-g}} (g_{\alpha\gamma} g_{\beta\delta} - g_{\alpha\delta} g_{\beta\gamma}) \quad (8)$$

where

$$m_{ik} m^{kj} \equiv \delta_i^j \quad (9)$$

and Greek indices are raised and lowered with the metric (2).

We observe that eq. (2) is invariant under $SL(3)$ transformations acting on the index i . It is natural to try to define a connection that "annihilates" the two-forms $\Sigma_{\alpha\beta i}$, in the sense that

$$\nabla_\alpha \Sigma_{\beta\gamma i} = \partial_\alpha \Sigma_{\beta\gamma i} - \Gamma_{\alpha\beta}{}^\delta \Sigma_{\delta\gamma i} - \Gamma_{\alpha\gamma}{}^\delta \Sigma_{\beta\delta i} + \mathcal{A}_{\alpha i}{}^j \Sigma_{\beta\gamma j}, \quad (10)$$

where $\Gamma_{\alpha\beta}{}^\gamma$ is a symmetric affine connection. Counting components, we find that this will admit a solution for the connection whenever the latter is an $SL(3)$ connection [3] ($SL(3)$ is eight-dimensional), and it may be verified that

the affine connection then becomes metric compatible, provided that the conformal factor σ is some function of the Σ 's. It is now straightforward to relate the $\text{SL}(3)$ curvature to the Riemann tensor of the metric (2):

$$0 = [\nabla_\alpha, \nabla_\beta] \Sigma_{\gamma\delta i} = R_{\alpha\beta\gamma}{}^\sigma \Sigma_{\sigma\delta i} - R_{\alpha\beta\delta}{}^\sigma \Sigma_{\sigma\gamma i} + \mathcal{F}_{\alpha\beta i}{}^j \Sigma_{\gamma\delta j}. \quad (11)$$

Here $\mathcal{F}_{\alpha\beta i}{}^j$ is the $\text{SL}(3)$ curvature tensor. It is easy to show that the symmetric part

$$\mathcal{F}_{\alpha\beta(ij)} = \mathcal{F}_{\alpha\beta(i}{}^k m_{j)k} = 0, \quad (12)$$

which means that the curvature lies in an $\text{SO}(3)$ subgroup of $\text{SL}(3)$. The connection may be written as

$$\mathcal{A}_{\alpha ij} \equiv \mathcal{A}_{\alpha i}{}^k m_{kj} \equiv i\epsilon_{ikj} \alpha_\alpha{}^k + \beta_{\alpha ij}; \quad (13)$$

$$\beta_{\alpha ij} = \frac{1}{2}(\partial_\alpha - \Gamma_{\alpha\gamma}{}^\gamma) m_{ij}. \quad (14)$$

Eq. (11) may be solved for the $\text{SL}(3)$ curvature, or alternatively for the self-dual part of the Riemann tensor, with the result that

$$R_{\alpha\beta\gamma\delta}^{(+)} = \frac{\sigma}{2} \epsilon^{ijk} m_{km} \mathcal{F}_{\alpha\beta j}{}^m \Sigma_{\gamma\delta i} \quad (15)$$

$$\mathcal{F}_{\alpha\beta i}{}^j = \frac{2i}{\sigma} \sqrt{-g} R_{\alpha\beta}{}^{\gamma\delta} \Sigma_{\gamma\delta k} \epsilon_{imn} m^{mj} m^{nk}. \quad (16)$$

Now we turn to dynamics. In writing down field equations, we prefer to use an $\text{SO}(3)$ connection, since the formalism simplifies. Moreover $\text{SO}(3)$ admits spinor representations while $\text{SL}(3)$ does not. So we switch to the formalism of ref. [2], and introduce an $\text{SO}(3)$ connection, its field strength, and the matrix

$$\Omega^{ij} \equiv \epsilon^{\alpha\beta\gamma\delta} F_{\alpha\beta}{}^i F_{\gamma\delta}{}^j. \quad (17)$$

Then we write an action of the general form [4]

$$S = \int \mathcal{L}(\eta; Tr\Omega, Tr\Omega^2, Tr\Omega^3) \quad (18)$$

where η is a Lagrange multiplier as well as a scalar density of weight minus one, and the traces are defined by means of the group metric of $SO(3)$, e.g.

$$Tr\Omega \equiv g_{ij}\Omega^{ij}. \quad (19)$$

In order to keep things reasonably specific, we will concentrate on the special case

$$S = \frac{1}{8} \int \eta (Tr\Omega^2 + \alpha (Tr\Omega)^2), \quad (20)$$

where α is a parameter. It is shown in ref. [2] that this action gives field equations that are equivalent to Einstein's equations for vanishing cosmological constant, provided that a certain non-degeneracy condition holds,

$$\alpha = -\frac{1}{2}, \quad (21)$$

and provided that the metric is defined by

$$g_{\alpha\beta} = \frac{8}{3} \eta \epsilon_{ijk} F_{\alpha\gamma}^i \tilde{F}^{\gamma\delta j} F_{\delta\beta}^k. \quad (22)$$

What about the general case? It is very important to observe that, an arbitrary conformal factor apart, the definition of the metric is not arbitrary. It is enforced by the requirement that the theory should have a Hamiltonian formulation which makes sense. This requirement also permits one to show that the same formula for the metric holds (up to conformal transformations) in the general case, so the previous equation is in fact a theorem. (For details, see ref. [4].) However, this metric is not Einstein unless $\alpha = -1/2$. The Hamiltonian formulation moreover shows that the number of degrees of freedom remains two per space-time point for all the theories that can be constructed in this way, which is why we refer to them as "neighbours of Einstein's equations". We observe that, by construction,

$$F_{\alpha\beta i} \in \mathbf{W}^+ \quad (23)$$

where $\mathbf{W}^+$ denotes the self-dual subspace.

Let us consider the field equations from the action (20):

$$\epsilon^{\alpha\beta\gamma\delta} D_\beta \Sigma_{\gamma\delta i} = 0 \quad (24)$$

$$\Sigma_{\alpha\beta i} \equiv \Psi_{ij} F_{\alpha\beta}^j; \quad \Psi_{ij} \equiv 2\eta(\Omega_{ij} + \alpha g_{ij} \text{Tr}\Omega) \quad (25)$$

$$\text{Tr}\Omega^2 + \alpha(\text{Tr}\Omega)^2 = 0. \quad (26)$$

In the second equation (which from now on is used to define Σ) we have, for the first and last time in this letter, used the group metric to lower indices. One can show that the choice of conformal factor in eq. (22) implies that

$$\sqrt{-g}g_{\alpha\beta} = \frac{i}{6}\epsilon^{ijk}(\Sigma_i \tilde{\Sigma}_j \Sigma_k)_{\alpha\beta}. \quad (27)$$

Hence the conformal factor of the metric can be expressed as a function of the Σ 's.

The first field equation may be used to solve for the connection in terms of the Σ 's. The calculation is straightforward and the result is important, but its explicit form is unilluminating, so we do not give it here. Of more immediate interest is the following consequence of the third equation (m_{ij} is defined in eq. (5)):

$$m_{ij} = 4\eta^2 \left((1 + 2\alpha)\Omega^2_{ij} \text{Tr}\Omega + (\alpha + \frac{1}{2})(\alpha - 1)\Omega_{ij}(\text{Tr}\Omega)^2 \right) - 8i\sqrt{-g}g_{ij}. \quad (28)$$

(Here $\sqrt{-g}$ denotes the square root of the space-time metric, not the group metric.) Hence the two natural metrics that exist on the $\text{SO}(3)$ fibers are proportional if and only if $\alpha = -1/2$, that is to say in the Einstein case. How are the $\text{SO}(3)$ and $\text{SL}(3)$ formalisms related? Introducing a metric on the fibers breaks $\text{SL}(3)$ down to $\text{SO}(3)$. However, in general we have two different fibre metrics available, m_{ij} and g_{ij} . Eq. (28) reveals that they are proportional only if $\alpha = -1/2$, that is to say in the Einstein case. In general we are dealing with two different $\text{SO}(3)$ subgroups. The $\text{SO}(3)$ field strength is self-dual by construction, and we see that

$$\mathcal{F}_{\alpha\beta i}{}^j \in \mathbf{W}^+ \quad \Leftrightarrow \quad m_{ij} \propto g_{ij}. \quad (29)$$

Referring back to eq. (15), we see that the self-dual "from the right" part of the Riemann tensor is self-dual "from the left" - a condition which implies the vanishing of the traceless part of the Ricci tensor - only when the two fibre metrics are proportional. We also see from eq. (28) that this is a quite exceptional case.

To add concreteness to the above, we will now consider a simple exact solution of eqs. (24-26). We choose the Ansatz [7]

$$A_{x1} = a_1(t) \quad A_{y2} = a_2(t) \quad A_{z3} = a_3(t), \quad (30)$$

all others zero ($A_{\alpha i}$ is the $\text{SO}(3)$ connection). If we choose the gauge

$$\eta = \frac{1}{16\dot{a}_1\dot{a}_2\dot{a}_3} \quad (31)$$

we find the solution

$$a_i = t^{-\frac{2c_i}{c}} \quad ; \quad c \equiv c_1 + c_2 + c_3, \quad (32)$$

where the integration constants have to obey

$$c_1^2 + c_2^2 + c_3^2 + \alpha c^2 = 0. \quad (33)$$

The resulting metric is

$$ds^2 = -dt^2 + t^{2\gamma_1} dx^2 + t^{2\gamma_2} dy^2 + t^{2\gamma_3} dz^2. \quad (34)$$

The Kasner exponents are defined by

$$\gamma_i \equiv 1 - \frac{2c_i}{c} \quad ; \quad \sum \gamma_i = 1 \quad \sum \gamma_i^2 = -1 - 4\alpha. \quad (35)$$

For $\alpha = -1/2$ this is the familiar Kasner metric. For other values of α the Ricci tensor turns out to have only one non-vanishing component, viz.

$$R_{tt} = 2(1 + 2\alpha)t^{-2}. \quad (36)$$

Finally, let us comment on the situation when the space-time dimension differs from four. It is unlikely that geometry can be based on two-forms in any dimension higher than four. In three dimensions, on the other hand, there is a natural isomorphism between the space of two-forms and the space of vectors, which we express explicitly through

$$*\Sigma^\alpha \equiv \frac{1}{2}\epsilon^{\alpha\beta\gamma}\Sigma_{\beta\gamma}. \quad (37)$$

Therefore, in a three dimensional space-time a "form connection" as defined by eq. (10) becomes just the ordinary $SO(1,2)$ spin connection. There is now no analogue of the four dimensional "neighbours" of Einstein's equations, since there is essentially only one object available from which the action may be constructed, namely $\det^*F$. Actually, this is not quite true, since a Chern-Simons term may be added. Consider

$$S = 2 \int \sqrt{\frac{1}{\lambda}} \det^* F + \theta \epsilon^{\alpha\beta\gamma} (A_{\alpha I} \partial_\beta A_\gamma^I + \frac{1}{3} f_{IJK} A_\alpha^I A_\beta^J A_\gamma^K). \quad (38)$$

The first term here gives Einstein's equations with $\lambda =$ the cosmological constant [8]. Including the second term gives a non-vanishing torsion tensor

$$T_{\alpha\beta}^I = 2\theta F_{\alpha\beta}^I. \quad (39)$$

However, the field strength may still be expressed in terms of a "metric" triad, so that the equation for the Ricci tensor eventually collapses to

$$R_{\alpha\beta} = 2\lambda(1 - \lambda\theta^2)g_{\alpha\beta}. \quad (40)$$

We see that the Chern-Simons term only rescales the cosmological constant. Its presence will be felt when matter fields are added.

Acknowledgements: This work was done while I was visiting Imperial College, London. I am grateful for the hospitality, and to the Science Research Council for financial support.

References

- [1] A. Ashtekar, Phys. Rev. **D36** (1987) 1587.
A. Ashtekar, Lectures on Non-Perturbative Canonical Gravity, World Scientific, Singapore 1991.
- [2] R. Capovilla, J. Dell and T. Jacobson, Phys. Rev. Lett. **63** (1989) 2325.
- [3] G. 'tHooft, Nucl. Phys. **B357** (1991) 211.
- [4] R. Capovilla, Nucl. Phys. **B373** (1992) 233
I. Bengtsson and P. Peldán, Int. J. Mod. Phys. **A7** (1992) 1287
I. Bengtsson, J. Math. Phys. **32** (1991) 3158
I. Bengtsson, Neighbours of Einstein's Equations - Some Further Results, Class. Quant. Grav., to appear.
- [5] J. Plebański, J. Math. Phys. **18** (1977) 2511
W. Israel, Differential Forms in General Relativity, Com. Dublin Inst. Adv. Stud. Ser. **A** No. 26, 1979.
L. Mason and J. Frauendiener, in R. Baston and T. Bailey (eds.): Twistors in Mathematical Physics, Cambridge U.P. 1990.
R. Capovilla, J. Dell, T. Jacobson and L. Mason, Class. Quant. Grav. **8** (1991) 41.
- [6] H. Urbantke, J. Math. Phys. **25** (1984) 2321
S.K. Donaldson and D.P. Sullivan, Acta Math. **163** (1990) 181.
G. Harnett, J. Phys. **A25** (1992) 5649.
- [7] R. Capovilla and J. Plebański, J. Math. Phys. **34** (1993) 130.
- [8] P. Peldán, Class. Quant. Grav. **9** (1992) 2079.